



## SCET sum rules for $B \rightarrow P$ and $B \rightarrow V$ transition form factors

Fulvia De Fazio  
INFN- Sezione di Bari

*Mainz, April 5th 2008*

### Outline

- heavy to light decays: - general considerations  
- comparison with heavy-to-heavy case
- SCET- based LCSR: hadronic form factors
- summary and perspectives

Based on work in collaboration with T. Feldmann and T. Hurth  
NPB B733 (2006) 1;  
JHEP02 (2008) 031

**Something well known to start with...**

## Heavy to light decays induced by $b \rightarrow u$ transition

$$B \rightarrow X_u \ell \nu$$

$$B \rightarrow \pi \ell \nu, \rho \ell \nu \dots$$



$$B \rightarrow \pi\pi, \rho\rho\dots$$

- relevant for the determination of  $V_{ub}$



**possibility to constrain one of the sides of the Unitarity Triangle**

- form factors describing exclusive semileptonic modes enter also in the description of some non leptonic decays in the QCD factorization approach



**possibility to constrain one of the angles of the Unitarity Triangle ( $\alpha$ )**

- description of rare B decays induced by  $b$  to  $s$  transition:

$$B \rightarrow K\pi, K^{(*)} \ell^+ \ell^-, K^* \gamma \dots$$

## Heavy-to-heavy decays: heavy quark symmetries

Give the possibility to relate the various form factors describing exclusive processes induced by  $b \rightarrow c$  transition

$$\langle D(v') | V_\mu | B(v) \rangle = \sqrt{m_B m_D} \left[ h_+(w) (v + v')_\mu + h_-(w) (v - v')_\mu \right],$$

$$\langle D^*(v', \epsilon') | V_\mu | B(v) \rangle = i\sqrt{m_B m_{D^*}} h_V(w) \epsilon_{\mu\nu\alpha\beta} \epsilon'^{\nu\alpha} v'^\alpha v^\beta,$$

$$\langle D^*(v', \epsilon') | A_\mu | B(v) \rangle = \sqrt{m_B m_{D^*}} \left[ h_{A_1}(w) (w + 1) \epsilon'_\mu - h_{A_2}(w) \epsilon'^* \cdot v v_\mu - h_{A_3}(w) \epsilon'^* \cdot v v'_\mu \right],$$

$$\langle D^*(v', \epsilon') | V_\mu | B^*(v, \epsilon) \rangle = \sqrt{m_{B^*} m_{D^*}} \left\{ -\epsilon \cdot \epsilon'^* \left[ h_1(w) (v + v')_\mu + h_2(w) (v - v')_\mu \right] + h_3(w) \epsilon'^* \cdot v \epsilon_\mu + h_4(w) \epsilon \cdot v' \epsilon'_\mu - \epsilon \cdot v' \epsilon'^* \cdot v \left[ h_5(w) v_\mu + h_6(w) v'_\mu \right] \right\},$$

$$\langle D^*(v', \epsilon') | A_\mu | B^*(v, \epsilon) \rangle = i\sqrt{m_{B^*} m_{D^*}} \epsilon_{\mu\nu\alpha\beta} \epsilon^\alpha \epsilon'^{\nu\beta} \left[ h_7(w) (v + v')^\nu + h_8(w) (v - v')^\nu \right]$$


At leading order in the heavy quark expansion one finds that

$$h_+ = h_V = h_{A_1} = h_{A_3} = h_1 = h_3 = h_4 = h_7 = \xi \quad \Rightarrow$$

all form factors are related to one **Isgur-Wise function**

## Heavy-to-heavy decays


In the kinematic regime when  $Q$  interacts with the light antiquark only through soft gluon exchanges heavy quark symmetries arise (**HQET**)



**HQET** reproduces correctly the long distance physics of **QCD**

Short-distance corrections, to be computed in perturbation theory

$$\alpha_s(m_Q) \approx 0.2$$



Hard vertex renormalization

## Heavy-to-light decays - Form factors

$$\langle P(p') | \bar{q} \gamma_\mu b | B(p) \rangle \rightarrow f_+(q^2), f_0(q^2)$$


$$\langle P(p') | \bar{q} \sigma_{\mu\nu} b | B(p) \rangle \rightarrow f_T(q^2)$$

$$\langle V(p', \varepsilon) | \bar{q} \gamma_\mu (1 - \gamma_5) b | B(p) \rangle \rightarrow V(q^2), A_1(q^2), A_2(q^2), A_0(q^2)$$

$$\langle V(p', \varepsilon) | \bar{q} \sigma_{\mu\nu} (1 - \gamma_5) b | B(p) \rangle \rightarrow T_1(q^2), T_2(q^2), T_3(q^2)$$

Are there symmetry relations also for heavy-to-light decays?

## Heavy-to-light form factors at large recoil



Large energy of the light meson  $E - \frac{M}{2} \ll M \iff q^2 \ll M^2$

If one assumes that the  $b$  and  $u$  still interact with the spectator only through “**soft**” gluon exchanges symmetry relations can be derived for the heavy-to-light form factors

Charles et al. PRD 60 (99) 014001

## Heavy-to-light form factors at large recoil

$$\left. \begin{aligned} \langle P(p') | \bar{q} \gamma_\mu b | B(p) \rangle \\ \langle P(p') | \bar{q} \sigma_{\mu\nu} b | B(p) \rangle \end{aligned} \right\} \longrightarrow \xi_P(E)$$

$$\left. \begin{aligned} \langle V(p', \varepsilon) | \bar{q} \gamma_\mu (1 - \gamma_5) b | B(p) \rangle \\ \langle V(p', \varepsilon) | \bar{q} \sigma_{\mu\nu} (1 - \gamma_5) b | B(p) \rangle \end{aligned} \right\} \longrightarrow \xi_\perp(E), \quad \xi_\parallel(E)$$

## Symmetry relations



$$f_+(q^2) = \frac{M}{2E} f_0(q^2) = \frac{M}{M + m_P} f_T(q^2) = \xi_P(E)$$

$$\frac{M}{M + m_V} V(q^2) = \frac{M + m_V}{2E} A_1(q^2) = T_1(q^2) = \frac{M}{2E} T_2(q^2) = \xi_\perp(E),$$



$$\frac{m_V}{E} A_0(q^2) = \frac{M + m_V}{2E} A_1(q^2) - \frac{M - m_V}{M} A_2(q^2) = \frac{M}{2E} T_2(q^2) - T_3(q^2) = \xi_\parallel(E)$$

## Symmetry relations

- Are valid for the soft contribution to the soft form factors at large recoil neglecting  $1/m_b$  and  $\alpha_s$  corrections
- Corrections stem from



Hard vertex  
renormalization




factorizable spectator  
interaction



The inclusion of such corrections leads to a ***factorization theorem***

# Energy scales




$$\Lambda_{QCD}$$

$$\mu_{hc} = \sqrt{m_b \Lambda}$$

$$m_b$$

**Soft** scale,  
typical of energies and momenta  
of light degrees of freedom



**Hard** scale

**Hard-collinear** scale

in interactions between soft and energetic modes  
in the initial and final state

## Separation of scales achieved in SCET

- $\mu < \mu_I$  SCET<sub>I</sub> (integrate out hard modes)
- $\mu < \mu_{II}$  SCET<sub>II</sub> (integrate out also hard-collinear modes)

Bauer, Fleming, Pirjol and Stewart  
Beneke, Chapovsky, Diehl and Feldmann  
Becher, Hill, Lange and Neubert  
...

# Factorization theorem

Beneke and Feldmann, NPB 685 (04) 249  
See also  
Hill and Neubert NPB 657 (03) 229  
Lange and Neubert NPB 690 (04) 249  
Bauer, Pirjol and Stewart PRD 67 (03) 071502

Short-distance functions  
Arise from integrating out hard modes:  $\mu_I < m_b$

Light-cone distributions

$$f_i(q^2) = C_i(E, \mu_I) \xi(E, \mu_I) + \underbrace{T_i(E, \mu_{II})}_{\text{Hard scattering kernel}} \otimes \Phi_B(\mu_{II}) \otimes \Phi_P(\mu_{II}) + \dots$$

Hard scattering kernel (perturbative series in  $\alpha_s$ )  
Contains the effects of both h and hc dynamics  $\mu_{II} < \mu_{hc}$

Soft form factor  
It is universal for each meson  
and does not depend on the Dirac structure  
of the current

Subleading terms in  $\Lambda / m_b$


The second term is the symmetry breaking one.  
The first enters already at tree level, while the second is  $O(\alpha_s)$ .  
However, the first could be Sudakov suppressed.

⇒ Their relative weight is a debated issue

# SCET – matching heavy-to-light currents

## Vector current

$$\bar{q}\gamma_\mu b \rightarrow C_3(\mu) [\bar{\chi}\gamma_\mu^\perp h_v] + \{C_4(\mu)n_{-\mu} + C_5(\mu)v_\mu\} [\bar{\chi}h_v]$$


$\chi = W\xi \rightarrow$  Hard-collinear light quark field  
 Wilson line

$$n_+^\mu = (1,0,0,-1)$$

$$n_-^\mu = (1,0,0,1)$$

At tree level  $C_3(m_b)=C_4(m_b)=1$   $C_5(m_b)=0$

The running is obtained solving RGEs in SCET



$$\langle M(p) | \bar{q}\Gamma b | B(v) \rangle \rightarrow \sum_i C_i(\mu) \langle M(p) | \bar{\chi}\Gamma_i h_v | B(v) \rangle + \Delta F_\Gamma$$

Wilson coeff. In SCET

Soft contribution

Hard, symmetry-breaking factorizable contribution

**B** → **P**

$$\langle P(p) | \bar{\chi} h_\nu | B(v) \rangle = 2E \xi(E)$$

$$\langle P(p) | \bar{\chi} \gamma_5 h_\nu | B(v) \rangle = 0$$

$$\langle P(p) | \bar{\chi} \gamma_\mu^\perp h_\nu | B(v) \rangle = 0$$

**B** → **V**

$$\langle V(p, \varepsilon) | \bar{\chi} h_\nu | B(v) \rangle = 0$$

$$\langle V(p, \varepsilon) | \bar{\chi} \gamma_5 h_\nu | B(v) \rangle = -2m_f \xi_{\parallel}(E) (v \cdot \varepsilon^*)$$

$$\langle V(p, \varepsilon) | \bar{\chi}_p \gamma_\mu^\perp h_\nu | B(v) \rangle = 2E \xi_{\perp}(E) i \varepsilon_{\perp}^{\mu\nu} \varepsilon_\nu^*$$

## Form factors

SCET provides a theoretical framework to achieve factorization of short and long distance physics.

However, non perturbative quantities should be determined through other theoretical approaches (lattice, QCD sum rules).

This is just the case of soft non-factorizable form factors

Other approaches (attempts to factorize the soft form factors):

- pQCD approach

Li, Mishima, Sanda

- zero-bin approach


Manohar, Stewart

## Light-cone QCD sum rules

Symmetry relations arise when the b quark decays to a highly energetic u quark and both interact with the spectator through “soft” gluon exchange

 soft contribution to the form factors

The two quarks in the final light meson are in an asymmetric configuration: one of them takes almost all the meson' momentum

 end-point of the wave function

In conventional light-cone sum rules soft contribution is related to terms proportional to the wave function at the end point (in the limit  $m_b \rightarrow \infty$  ).

 a-posteriori identification of the soft form factors

## Sum rules in SCET

The starting point is a correlation function:

The external state is the heavy meson in contrast to usual LCsr

$$\Pi(p') = i \int d^4x \ e^{ip' \cdot x} \ \langle 0 | T [ J_M(x) J_\Gamma(0) ] | B(p_B = m_B \mathbf{v}) \rangle$$

Interpolating current for the light meson

Current inducing the transition

We work in the frame:

$$p'_\perp = v_\perp = 0 \quad n_+ \cdot v = n_- \cdot v = 1$$

2 independent variables

$$\Rightarrow \quad n_+ \cdot p' \cong 2E = O(m_b) \quad 0 > n_- \cdot p' \cong O(\Lambda)$$

we work for fixed values of  $(n_+ \cdot p')$ , so that

$$\Pi = \Pi(n_- \cdot p')$$

**B**  $\rightarrow$   $\pi$

$$J_M(x) = J_\pi(x) = -i \bar{\chi}(x) \not{n}_+ \gamma_5 \chi(x) - i (\bar{\chi}(x) \not{n}_+ \gamma_5 q_s(x) + h.c.)$$

$$J_\Gamma(0) = J_0(0) = \bar{\chi}(0) h_v(0)$$

**B** → **V**

Correlators:

$$\Pi_{\parallel}(n_{-} \cdot p') = i \int d^4x e^{ip' \cdot x} \langle 0 | T [J_V^{\parallel}(x) J_0^{\parallel}(0)] | B(v) \rangle$$

$$\frac{1}{2} \varepsilon^{\mu_1 \nu_1 \sigma \tau} n_{+\sigma} n_{-\tau} \Pi_{\perp}(n_{-} \cdot p') = i \int d^4x e^{ip' \cdot x} \langle 0 | T [J_V^{\mu_{\perp}}(x) J_0^{\nu_{\perp}}(0)] | B(v) \rangle$$

Currents

$$J_V^{\parallel}(x) = -i \bar{\chi}(x) \not{n}_+ \chi(x) - i (\bar{\chi}(x) \not{n}_+ q_s(x) + h.c.)$$

$$i J_V^{\mu_{\perp}}(x) = \bar{\chi}(x) i \not{n}_+ \gamma^{\mu_{\perp}} \chi(x) + (\bar{\chi}(x) i \not{n}_+ \gamma^{\mu_{\perp}} q_s(x) + h.c.)$$

$$J_0^{\parallel} = \bar{\chi}(-\gamma_5) h_v$$

$$J_0^{\nu_{\perp}} = \bar{\chi}(\gamma^{\nu_{\perp}}) h_v$$

Matrix elements

$$\langle 0 | J_V^{\parallel} | V(p', \varepsilon) \rangle = m_V (n_+ \cdot \varepsilon) f_V^{\parallel}$$

$$\langle 0 | i J_V^{\mu_{\perp}} | V(p', \varepsilon) \rangle = (n_+ \cdot p') \varepsilon^{\mu_{\perp}} f_V^{\perp}$$

## Tree level sum rule: $B \longrightarrow \pi$

The procedure consists in writing the correlator in two different ways (as with all QCD sum rule calculations)


Hadronic side

$$\Pi^{\text{HAD}}(n_- p') = \Pi(n_- p') \Big|_{\text{res.}} + \Pi(n_- p') \Big|_{\text{cont.}}$$

↓  
contribution of the pion


↘  
Higher states and continuum;  
starts above an effective threshold  $\omega_s$

Using  $\langle 0 | J_\pi | \pi(p') \rangle = (n_+ \cdot p') f_\pi \quad \langle \pi(p') | J_0(0) | B(m_B v) \rangle = (n_+ \cdot p') \xi_\pi(n_+ \cdot p', \mu_I)$



$$\Pi(n_- \cdot p') \Big|_{\text{res}} = - \frac{(n_+ \cdot p') \xi_\pi(n_+ \cdot p') f_\pi}{n_- \cdot p'}$$

## SCET side



Hard-collinear quark propagator

$$S_F^{\text{hc}} = \frac{i}{n_- p' - \omega + i\eta} \frac{\not{n}_-}{2}$$

$$\omega = n_- \cdot k$$

$k^\mu$  being the momentum of the soft light quark that ends up as spectator in the B

Result:

$$\Pi(n_- p') = f_B m_B \int_0^\infty d\omega \frac{\phi_-^B(\omega)}{\omega - n_- p' - i\eta}$$

where the B light-cone distribution amplitude enters through

$$\langle 0 | \bar{q}_s(x_-) \gamma_5 \frac{\not{n}_+ \not{n}_-}{2} h_v(0) | B(p_B) \rangle = -i f_B m_B \int d\omega' e^{-i\omega' \frac{n_+ \cdot x}{2}} \phi_-^B(\omega')$$

This has already the form of a dispersion relation in  $n_- \cdot p'$ :

$$\Pi(n_- p') = \frac{1}{\pi} \int_0^\infty d\omega' \frac{\text{Im}[\Pi(\omega')]}{\omega' - n_- p' - i\eta}$$

with 
$$\frac{1}{\pi} \text{Im}[\Pi(\omega')] = f_B m_B \phi_-^B(\omega')$$

Also  $\Pi(n_- \cdot p')|_{cont.}$  can be written as a dispersion relation where, assuming global quark-hadron duality, we identify the spectral density with the expression obtained in SCET

$$\Pi(n_- \cdot p')|_{cont.} = f_B m_B \int_{\omega_s}^{\infty} d\omega \frac{\phi_-^B(\omega)}{\omega - n_- \cdot p' - i\eta}$$

Sum rule obtained equating the two representations:

$$-\frac{(n_+ \cdot p') \xi_\pi(n_+ \cdot p') f_\pi}{n_- \cdot p'} = f_B m_B \int_0^{\omega_s} d\omega \frac{\phi_-^B(\omega)}{\omega - n_- \cdot p' - i\eta}$$

Borel transformation:

$$\hat{B}(\omega_M) \frac{1}{\omega - n_- \cdot p'} = \frac{1}{\omega_M} e^{-\omega/\omega_M}$$

**Final Sum Rule – tree level**

$$\xi_\pi(n_+ \cdot p') = \frac{f_B m_B}{(n_+ \cdot p') f_\pi} \int_0^{\omega_s} d\omega e^{-\omega/\omega_M} \phi_-^B(\omega)$$


Tree level sum rule:  $B \rightarrow V_{\parallel}, V_{\perp}$

$$\begin{aligned}\hat{\xi}_{\parallel}(n+p') &\equiv \frac{n+p'}{2m_V} \xi_{\parallel}(n+p') \\ &= \frac{f_B m_B}{f_V^{\parallel}(n+p')} \exp\left[\frac{m_V^2}{(n+p')\omega_M}\right] \int_0^{\omega_s} d\omega' e^{-\omega'/\omega_M} \phi_B^-(\omega')\end{aligned}$$

Identical to the pion case except that now  $m_V^2 \neq 0$ ,  $f_{\pi} \rightarrow f_V^{\parallel}$

$$\xi_{\perp}(n+p') = \frac{f_B m_B}{f_V^{\perp}(n+p')} \exp\left[\frac{m_V^2}{(n+p')\omega_M}\right] \int_0^{\omega_s} d\omega' e^{-\omega'/\omega_M} \phi_B^-(\omega')$$

# Radiative corrections from hard-collinear loops



$$\Pi(n_- \cdot p') = f_B m_B \int_0^\infty d\omega \frac{\phi_-^B(\omega)}{\omega - n_- \cdot p' - i\eta} \left\{ 1 + \frac{\alpha_s C_F}{4\pi} [(a1 - a4) + (b1 - b2)] \right\}$$

## Results

$$\hat{\xi}_{\parallel}(\xi_{\perp}) = \frac{f_B m_B}{f_V^{\parallel(\perp)} (n+p')} \exp \left[ \frac{m_V^2}{(n+p') \omega_M} \right] \int_0^{\omega_s} d\omega' e^{-\omega'/\omega_M} \phi_{\parallel(\perp)}^{\text{eff}}(\omega')$$

$$\phi_{\parallel,\perp}^{\text{eff}}(\omega', n+p', \mu) \equiv \left\{ - \int_{\omega'}^{\infty} d\omega f_{\parallel,\perp}(\omega, \omega', \mu) \frac{d\phi_B^-(\omega, \mu)}{d\omega} + \int_0^{\omega'} d\omega \left[ \frac{g_{\parallel,\perp}(\omega, \omega', \mu)}{\omega - \omega'} \right]_+ \phi_B^-(\omega, \mu) \right\}$$



$\mu$ -dependence through  $L_0 = \ln \left[ \frac{\mu^2}{(n+p') \omega'} \right]$

## Renormalization scale dependence

The renormalization scale dependence of the form factors should cancel against that of the Wilson coefficients:

$$\frac{d}{d \ln \mu} C_i(\mu) = -\frac{\alpha_s C_F}{4\pi} \left( \Gamma_{\text{cusp}}^{(1)} \ln \frac{\mu}{m_b} + 5 \right) C_i(\mu) + \dots$$

This can be checked knowing the scale dependence of  $\phi_B^-$

$$\frac{d}{d \ln \mu} \phi_B^-(\omega; \mu) = -\frac{\alpha_s C_F}{4\pi} \int_0^\infty d\tilde{\omega} \gamma_-^{(1)}(\omega, \tilde{\omega}; \mu) \phi_B^-(\tilde{\omega}; \mu) + \dots$$

↓  
anomalous dimension

$$\begin{aligned} \gamma_-^{(1)}(\omega, \tilde{\omega}; \mu) = & \left( \Gamma_{\text{cusp}}^{(1)} \ln \frac{\mu}{\omega} - 2 \right) \delta(\omega - \tilde{\omega}) - \Gamma_{\text{cusp}}^{(1)} \frac{\theta(\tilde{\omega} - \omega)}{\tilde{\omega}} \\ & - \Gamma_{\text{cusp}}^{(1)} \omega \left[ \frac{\theta(\tilde{\omega} - \omega)}{\tilde{\omega}(\tilde{\omega} - \omega)} \right]_+ - \Gamma_{\text{cusp}}^{(1)} \omega \left[ \frac{\theta(\omega - \tilde{\omega})}{\omega(\omega - \tilde{\omega})} \right]_+ \end{aligned}$$

→ T. Feldmann, G. Bell  
arXiv:0711.4014




Complete cancellation of  $\mu$ -dependent terms

# Numerical results I: Soft form factor for longitudinal vector mesons

**V=ρ**

$$n_+ \cdot p' = m_B$$

$$\mu = 1 \text{ GeV}$$



$$\Rightarrow \omega_s = \frac{m_S^2}{m_B^2} = \{0.35, 0.4, 0.45\}$$

next vector resonance being  $\rho(1450)$



Stability requirement:

$$D = \frac{\omega_M}{\xi_{||}} \frac{\partial \xi_{||}}{\partial \omega_M} < 25\%$$


Lower bound

Further constraint:  
continuum contribution not too large:

$$R = \frac{\int_0^{\omega_s} d\omega' e^{-\omega'/\omega_M} \phi_{||}^{eff}(\omega')}{\int_0^{\infty} d\omega' e^{-\omega'/\omega_M} \phi_{||}^{eff}(\omega')} > 50\%$$

Upper bound  
on  $\omega_M$

## Numerical results I: Soft form factor for longitudinal vector mesons

$$\hat{\xi}_{\parallel}(n_+ \cdot p' = m_B) = 0.33 \pm_{0.02}^{0.02} \Big|_{\omega_s} \pm_{0.06}^{0.03} \Big|_{\omega_M} \pm_{0.02}^{0.03} \Big|_{\omega_0} \pm 0.05 \Big|_{f_B}$$


dependence on the B meson wave function

$$\phi_B^-(\omega) = \frac{1}{\omega_0} e^{-\omega/\omega_0} \quad \omega_0(1 \text{ GeV}) = (0.48 \pm 0.05) \text{ GeV}$$

Lee, Neubert

Adding errors in quadrature:

$$\hat{\xi}_{\parallel}(n_+ \cdot p' = m_B) = 0.33 \pm_{0.09}^{0.07}$$




Compatible with the value obtained from traditional sum rules

Ball, Braun  
Ball, Zwicky  
Beneke, Feldmann, Seidel

# Numerical results I: Soft form factor for longitudinal vector mesons

## Energy dependence

----- Tree level



Fitted curve:

$$\hat{\xi}_{\parallel}(n_+ \cdot p') = \hat{\xi}_{\parallel}(m_B) \left( -a + \frac{b}{(n_+ \cdot p')} + c (n_+ \cdot p') \right)$$


$$a = 1.92 \pm_{0.17}^{0.42} \quad b = 12.25 \pm_{1.12}^{2.65} \quad c = 0.114 \pm_{0.01}^{0.024}$$

→ Including only the uncertainty on  $\omega_M$

# Numerical results II: Soft form factor for transverse vector mesons

$$n_+ \cdot p' = m_B$$

**V=ρ**



$$\longrightarrow \omega_s = \{0.20, 0.25, 0.30\}$$

Lower values in order  
to get rid of  $b_1(1235)$

$$\xi_{\perp}(n_+ \cdot p' = m_B) = 0.26 \pm_{0.04}^{0.03} \Big|_{\omega_s} \pm_{0.01}^{0.01} \Big|_{\omega_M} \pm 0.03 \Big|_{\omega_0} \pm 0.04 \Big|_{f_B}$$




$$\xi_{\perp}(n_+ \cdot p' = m_B) = 0.26 \pm_{0.07}^{0.06}$$

# Numerical results II: Soft form factor for transverse vector mesons

## Energy dependence

----- Tree level



Fitted curve:

$$\xi_{\perp}(n_{+} \cdot p') = \xi_{\perp}(m_B) \left( -a + \frac{b}{(n_{+} \cdot p')} + \frac{c}{(n_{+} \cdot p')^2} \right)$$


$$a = 0.26 \pm_{0.15}^{0.03} \quad b = 2.49 \pm_{2.63}^{0.69} \quad c = 21.76 \pm_{2.58}^{9.81}$$

→ Including only the uncertainty on  $\omega_M$

# Numerical results III: Soft form factor for pseudoscalar mesons

$$n_+ \cdot p' = m_B$$

**P= $\pi$**



$$\longrightarrow \omega_s = \{0.15, 0.20, 0.25\}$$

$$\xi_\pi(n_+ \cdot p' = m_B) = 0.25 \pm_{0.06}^{0.05} \Big|_{\omega_s} \pm_{0.03}^{0.02} \Big|_{\omega_M} \pm_{0.02}^{0.03} \Big|_{\omega_0} \pm_{f_B} 0.04$$




$$\xi_\pi(n_+ \cdot p' = m_B) = 0.25 \pm_{0.08}^{0.07}$$

# Numerical results III: Soft form factor for pseudoscalar mesons

## Energy dependence

----- Tree level



Fitted curve:

$$\xi_\pi(n_+ \cdot p') = \xi_\pi(m_B) \left( -a + \frac{b}{(n_+ \cdot p')} + \frac{c}{(n_+ \cdot p')^2} \right)$$

$$a = 0.23 \pm_{0.08}^{0.06}$$




$$b = 2.53 \pm_{1.27}^{0.94}$$

$$c = 21.00 \pm_{3.46}^{4.71}$$

→ Including only the uncertainty on  $\omega_S$

## Numerical results : modifying the sum rules

Possibility to decrease the sensitivity to sum rule parameters:  
Dividing for the sum rules relative to the decay constants



Reduction of the uncertainties  
due to  $\omega_s, \omega_M$   
but introduction of an unknown  
systematic error:  
Are the sum rules really correlated?

# Dependence on the shape of the B meson wave function



Default model:

$$\phi_B^-(\omega) = \frac{1}{\omega_0} e^{-\omega/\omega_0} \quad \omega_0(1 \text{ GeV}) = (0.48 \pm 0.05) \text{ GeV}$$

Alternatives:

$$\phi_B^-(\omega) = \frac{1}{\omega_0} \exp \left[ - \left( \frac{\omega}{\omega_1} \right)^2 \right], \quad \omega_1 = \frac{2\omega_0}{\sqrt{\pi}};$$

$$\phi_B^-(\omega) = \frac{1}{\omega_0} \left( 1 - \sqrt{\left( 2 - \frac{\omega}{\omega_2} \right) \frac{\omega}{\omega_2}} \right) \theta(\omega_2 - \omega), \quad \omega_2 = \frac{4\omega_0}{4 - \pi}$$




Sizable dependence  
More information on  $\phi_B^-$  required

## Form factor ratios


ratio:	$(\xi_\pi/f_\pi) : (\hat{\xi}_\rho^\parallel/f_\rho^\parallel)$	$(\xi_\rho^\perp/f_\rho^\perp) : (\hat{\xi}_\rho^\parallel/f_\rho^\parallel)$
original sum rule	$1.18^{+0.37}_{-0.32}$	$1.02^{+0.28}_{-0.21}$
modified sum rule	$1.05^{+0.06}_{-0.04} \pm (?)_{\text{syst.}}$	$0.87^{+0.06}_{-0.12} \pm (?)_{\text{syst.}}$

# Symmetry breaking corrections

Stem from:



Hard vertex  
renormalization



Hard-collinear spectator  
interaction

Explicit calculation gives

Beneke and Feldmann, NPB 592 (01) 3

$$\frac{f_0}{f_+} = \frac{2E}{M} \left( 1 + \frac{\alpha_s C_F}{4\pi} [2 - 2L] + \frac{\alpha_s C_F}{4\pi} \frac{M(M - 2E)}{(2E)^2} \frac{\Delta F_\pi}{\xi_\pi} \right)$$

Where  $\Delta F_\pi$  parametrizes the factorizable form factor contribution:

$$\Delta F_\pi = \frac{8\pi^2 f_B f_\pi}{N_C M} \left( \int dl_+ \frac{\phi_+^B(l_+)}{l_+} \right) \left( \int du \frac{\phi_\pi(u)}{1-u} \right)$$

## Factorizable form factor

$$\Pi(p') = i \int d^4x e^{ip' \cdot x} \langle 0 | T [J_\pi(x) J_1(0)] | B(p_B) \rangle \quad J_1 = \bar{\chi} g A_{hc}^\perp h_\nu$$

### Hadronic side

$$\hat{B}[\Pi_1](\omega_M) |_{had} = -\frac{\alpha_s C_F}{4\pi} \frac{f_\pi m_B^2}{2\omega_M} \Delta F_\pi e^{-m_\pi^2 / (n_+ \cdot p') \omega_M}$$

↓  
gives access to the factorizable term

The final sum rule has the same structure in the three cases:

$$\Delta F_X(\mu, n_+ p') = \frac{2f_B \omega_M (n_+ p')}{m_B f_X} e^{m_X^2 / (n_+ p' \omega_M)} \times \int_0^\infty \frac{d\omega}{\omega} \phi_B^+(\omega, \mu) \left( 1 - e^{-\omega_s / \omega_M} \theta(\omega - \omega_s) - e^{-\omega / \omega_M} \theta(\omega_s - \omega) \right)$$

with  $X = \pi, \rho_\parallel, \rho_\perp$

the parameters depending on each of the three cases

## Factorizable form factor

Inserting the leading order sum rule for the decay constants:

$$4\pi^2 f_X^2 \simeq M^2 e^{m_X^2/M^2} \left(1 - e^{-s_0/M^2}\right)$$

$$M^2 = \omega_M(n_+ \cdot p'), \quad s_0 = \omega_s(n_+ \cdot p')$$






$$\Delta F_X(\mu, n_+ p') = \frac{8\pi^2 f_B f_X}{m_B} \int_0^\infty \frac{d\omega}{\omega} \phi_B^+(\omega, \mu) \times \\ \times \frac{1 - e^{-\frac{s_0}{M^2}} \theta\left(\omega - \frac{s_0}{n_+ p'}\right) - e^{-\omega n_+ p'/M^2} \theta\left(\frac{s_0}{n_+ p'} - \omega\right)}{1 - e^{-\frac{s_0}{M^2}}}$$

In the limit  $s_0 \ll \omega(n_+ \cdot p')$  the QCD factorization result is recovered

$$\Delta F_X(\mu) \Big|_{\text{QCDF, asympt.}} = \frac{8\pi^2 f_B f_X}{m_B} \int_0^\infty \frac{d\omega}{\omega} \phi_B^+(\omega, \mu)$$

# Corrections to symmetry relations: numerics

original sum rule  
modified sum rule



- the dependence on  $\omega_M$  is milder for the modified sum rule
- qualitative agreement at this level of accuracy with the QCDF result i.e. no enhancement of the symmetry breaking term

## Conclusions

- light cone sum rules for exclusive B decays at large recoil can be derived within SCET
- non perturbative ingredients:
  - light cone wave functions of the B meson
  - sum rule parameters
- outcome consistent with full QCD results
- main result: possibility to separate from the very beginning soft - non factorizable term from the factorizable form factor
- hierarchy among soft/hard contributions in agreement with QCD factorization

At next to leading order

$$f_+(q^2) = \xi(E) \left[ C_4 + \frac{E}{m_B} C_5 \right]$$

$$f_0(q^2) = \xi(E) 2 \frac{E}{m_B} \left[ C_4 + \left( 1 - \frac{E}{m_B} \right) C_5 \right]$$

$$f_T(q^2) = \xi(E) C_{11}$$

$$A_1(q^2) = \xi_{\perp}(E) 2 \frac{E}{m_B} C_3$$

$$A_2(q^2) = \xi_{\perp}(E) C_3$$

$$V(q^2) = \xi_{\perp}(E) C_3$$

$$A_0(q^2) = \xi_{\parallel}(E) \left[ C_4 + C_5 \left( 1 - \frac{E}{m_B} \right) \right]$$

$$T_1(q^2) = \xi_{\perp}(E) C_9$$

$$T_2(q^2) = \xi_{\perp}(E) 2 \frac{E}{m_B} C_9$$

$$T_3(q^2) = \xi_{\perp}(E) C_9$$

## Factorizable form factor: $B \rightarrow \pi$ case

The spectator scattering terms can be identified by comparing  $B \rightarrow \pi$  form factors for different Dirac structures

Tree level matching (in the light-cone gauge) gives:


$$\bar{\psi}Q \rightarrow J_0 - \frac{1}{n_+ \cdot p'} J_1 \quad \bar{\psi}h_+Q \rightarrow 2J_0 - \frac{2}{m_b} J_1$$

$$J_0 = \bar{\chi}h_\nu \quad J_1 = \bar{\chi}gA_{hc}^\perp h_\nu$$

Using:  $\langle \pi(p') | \bar{q} \gamma_\mu b | B(p) \rangle = f_+(q^2) \left[ p_\mu + p'_\mu - \frac{m_B^2}{q^2} q_\mu \right] + f_0(q^2) \frac{m_B^2}{q^2} q_\mu$

$$\langle \pi(p') | \bar{q} b | B(p) \rangle = \frac{m_B^2}{m_b} f_0(q^2)$$

and  $q^2 = m_B(m_B - n_+ \cdot p')$



$$m_B f_0 = \langle \pi | J_0 | B \rangle - \frac{1}{n_+ p'} \langle \pi | J_1 | B \rangle,$$


$$(n_+ p') f_+ + m_B f_0 = 2 \langle \pi | J_0 | B \rangle - \frac{2}{m_B} \langle \pi | J_1 | B \rangle.$$




$$f_0/f_+ = \frac{n_+ p'}{m_B} \left( 1 - \frac{2q^2}{m_B^3} \frac{\langle \pi | J_1 | B \rangle}{\langle \pi | J_0 | B \rangle} + \mathcal{O}(\alpha_s^2) \right)$$

## Factorizable form factor

Comparing the two expressions for the ratio  $\frac{f_0}{f_+}$



$$\frac{\alpha_s C_F}{4\pi} \Delta F_\pi = -\frac{2}{m_B^2} \langle \pi | J_1 | B \rangle + \dots$$



In SCET-QCD-sum rules we need to consider a correlator involving the current  $J_1$